



BMJ Open Fitness, strength and severity of COVID-19: a prospective register study of 1 559 187 Swedish conscripts

Agnes af Geijerstam ,¹ Kirsten Mehlig,¹ Mats Börjesson,^{2,3} Josefina Robertson,^{1,4} Jenny Nyberg ,⁵ Martin Adiels,^{1,2} Annika Rosengren,^{2,6} Maria Åberg,^{1,7} Lauren Lissner¹

To cite: af Geijerstam A, Mehlig K, Börjesson M, *et al.* Fitness, strength and severity of COVID-19: a prospective register study of 1 559 187 Swedish conscripts. *BMJ Open* 2021;**11**:e051316. doi:10.1136/bmjopen-2021-051316

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-051316>).

Received 15 March 2021
Accepted 16 June 2021

ABSTRACT

Objective To investigate the possible connection between cardiorespiratory fitness (CRF) and muscle strength in early adulthood and severity of COVID-19 later in life.

Design Prospective registry-based cohort study.

Participants 1 559 187 Swedish men, undergoing military conscription between 1968 and 2005 at a mean age of 18.3 (SD 0.73) years.

Main outcome measures Hospitalisation, intensive care or death due to COVID-19 from March to September 2020, in relation to CRF and muscle strength.

Results High CRF in late adolescence and early adulthood had a protective association with severe COVID-19 later in life with OR (95% CI) 0.76 (0.67 to 0.85) for hospitalisation (n=2 006), 0.61 (0.48 to 0.78) for intensive care (n=445) and 0.56 (0.37 to 0.85) for mortality (n=149), compared with the lowest category of CRF. The association remains unchanged when controlled for body mass index (BMI), blood pressure, chronic diseases and parental education level at baseline, and incident cardiovascular disease before 2020. Moreover, lower muscle strength in late adolescence showed a linear association with a higher risk of all three outcomes when controlled for BMI and height.

Conclusions Physical fitness at a young age is associated with severity of COVID-19 many years later. This underscores the necessity to increase the general physical fitness of the population to offer protection against future viral pandemics.

BACKGROUND

The COVID-19 pandemic has affected the world in unprecedented ways. While measures such as contact restrictions and vaccinations relieve the immediate impact of the disease, epidemiological studies may identify risk factors that could be managed in the long run, in order to reduce the impact of future epidemics. Early on, advanced age and current cardiovascular comorbidities including obesity were found to be associated with a more severe course of COVID-19.¹⁻³

Cardiorespiratory fitness (CRF) is regarded as one of the most important risk factors for overall health, cardiovascular morbidity and overall mortality.⁴ The level of physical

Strengths and limitations of this study

- Data from the Swedish military service conscription registry provided us with objective measures of fitness in a uniquely large sample.
- The prospective design with long and near-complete follow-up from well-validated hospital records and the death registry gives a strong ground for drawing conclusions.
- Limitations include that it was not possible to distinguish conscripts with low versus extremely low fitness levels.
- Since female conscription was voluntary, our results can only be generalized to men.

activity early in life (and associated CRF) may have short-term and long-term effects on risk factors and the immune system. There has been speculation about the way it may confer protection against severe COVID-19.⁵ An early observation from the UK Biobank cohort that slower self-reported walking pace predicted more severe COVID-19 suggests the importance of CRF for this particular disease,⁶ and more recent studies have reinforced this notion, using self-reported measures^{7 8} but also clinically tested CRF and strength.^{9 10}

Although CRF is a persistent trait with a genetic component,¹¹ it is also to a large extent modifiable by regular exercise. Overall, physical fitness consists of CRF and muscular fitness, including muscle strength and healthy weight. Body mass index (BMI) is routinely measured in hospital and public care settings and has already been shown to be associated with severity of disease.^{2 12} However, objective measures of CRF and strength before the onset of COVID-19 are rarely available in population-based studies. Self-reported measures are unreliable, and most studies conducted in middle-aged or older populations have limited information on risk factors in earlier life.¹³ Participation rates among



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Agnes af Geijerstam;
agnes.af.geijerstam@gu.se

young men in population-based studies also tend to be low.¹⁴

Swedish military service conscription registry contains detailed and high-quality information about CRF and strength in a uniquely large sample of young men. In contrast to many population-based surveys, the participation rates until the early 2000s suggested highly generalisable results. Combined with measured data on BMI and other potential confounding factors in early adulthood, linkage data from nationwide disease registries enable a life course perspective on the role of physical fitness as a potentially mitigating factor for severe COVID-19 that is presently lacking in the literature.

METHODS

Study design

This prospective cohort study is based on data from the Swedish military service conscription registry, the Longitudinal integration database for health insurance and labour market studies (LISA), and the Swedish national hospital, intensive care and cause of death registries (for description in detail, see Åberg *et al.*).¹⁵ Fitness data were collected in early adulthood, in relation to incidence data on COVID-19 retrieved in September 2020, after the first wave of the pandemic in Sweden.

Patient and public involvement statement

As this is a registry-based study, there has been no patient or public involvement.

Population studied

The Swedish military service conscription registry contains information about 1 949 891 Swedish individuals who enlisted for military service between late 1968 and 2005 giving between 15 and 52 years of follow-up until the outbreak of COVID-19. During that time, Swedish law required all male citizens to enlist, except for those in prison or those with severe chronic somatic or psychiatric conditions or functional disabilities (approximately 2%–3% annually). Due to a shift to voluntary recruitment in 2005, the data in later years is no longer considered representative. The standardised protocol included measurements of weight and height, blood pressure, muscular strength and CRF. The registry also includes prior medical diagnoses. The examinations took place over 2 days in six conscription centres across Sweden.

Main independent variables

CRF score

To evaluate CRF, the subjects performed a cycle ergometric test where the work rate was successively increased until limited by exhaustion. Based on the final work rate, a nine-level score derived by the National Service Administration in Sweden¹⁶ served as a standardised measure of CRF. These scores were validated in relation to different tasks the conscripts were expected to manage during their time in the military (cross-country running with a

heavy backpack and heavy lifting).¹⁶ Over the decades, there have been some changes in the conscription examination protocols, and raw data were not available for all conscripts.¹⁷ However, the CRF scores matching certain military requirements were assigned during all recruitment years and can be assumed to be comparable across years.¹¹ For this study, the scores were collapsed into three categories describing low,^{1–5} medium^{6–7} and high^{8–9} CRF. The asymmetrical assignment of CRF scores was prompted by the fact that the fitness data in the lowest categories were omitted from the registry after 1999.¹⁵

Muscle strength score

Isometric muscle strength in units of newton was assessed as a weighted sum of knee extension (weight 1.3), elbow flexion (weight 0.8) and handgrip (weight 1.7). As the methods for measuring muscle strength did not change over the years and both the weighted values and assigned strength scores were available in the registry until 1996, it was possible to examine strength as a continuous variable.¹⁶

OTHER COVARIATES

BMI, height and weight status

Weight and height were measured by standard anthropometric measurement techniques, and continuous BMI values (kg/m^2) were calculated and also divided into categories of underweight (BMI<18.5), normal weight (BMI 18.5–24.9), overweight (BMI 25.0–29.9), obesity (BMI 30–34.9) and obesity class 2 (BMI≥35).¹⁸ For those aged 15–17 years at conscription, weight status categories were adjusted based on age-specific and sex-specific BMI z-scores, according to the WHO recommended childhood BMI cutoffs.¹⁹ In addition, height was included as a covariate, given its positive association with muscular strength.

Blood pressure

High blood pressure in early adulthood was used as an indicator of predisposition to later hypertension and other cardiovascular diseases (CVDs).⁶ Systolic and diastolic blood pressure was measured according to a standardised protocol after 5–10 min of rest in the supine position. In this study, values were then categorised into five subgroups (optimal blood pressure (systolic blood pressure <120 mm Hg and diastolic blood pressure <80 mm Hg), normal blood pressure (120–129 and 80–84 mm Hg), high normal blood pressure (130–139 and 85–89 mm Hg), grade 1 hypertension (140–159 and 90–99 mm Hg) and grade 2 hypertension (≥160 and ≥100 mm Hg) based on the 2018 European guidelines.²⁰

Chronic disease at baseline

To assess the specific effect of physical fitness, we controlled for morbidity at baseline, using International Classification of Diseases (ICD) codes for respiratory disease




Figure 1 Construction of analytic sample. BMI, body mass index; CRF, cardiorespiratory fitness.

(J00-J99), CVD (I00-I99), diabetes (E10-I4), kidney disease (N00-N29) and malignant cancers (C00-C97).

Parental education

Parental education was considered a proxy for socioeconomic position of the family of origin. Using data from the LISA registry, parental education was based on the highest of maternal and paternal education and divided into three categories with low education (up to 9 years), medium education (high school diploma with ≤ 2 years at university) and high (≥ 3 years at university).¹⁷

Analytic samples and subsamples

Figure 1 describes the construction of the analytic sample. Among 1 949 891 individuals examined for conscription between 1968 and 2005, the following exclusions were made: those with ambiguous identification numbers or without known place or year of conscription, those dying or emigrating without return before 2020 and all female conscripts, leaving 1 737 217 observations. Because of the aim of relating fitness measured in adolescence and early adulthood to disease severity later in life, individuals with age at conscription outside the interval of 15–30 years were excluded. Since BMI was an important confounder for the association between fitness and disease, the sample was further restricted excluding all men with missing or implausible BMI values (range for inclusion set between 15 and 60). Among the remaining 1 559 187 men, 1 143 670 had values for CRF and 1 161 914 had registered values for the continuous measure of muscle strength.

Outcome parameters: hospitalisation and/or death due to COVID-19

Using the Swedish personal identification number, the full sample was linked to the national hospital, the intensive care and the cause of death registries. From these, all cases between March and September 2020 with a main diagnosis of ICD U071 for test verified infection with

SARS-CoV-2 and U072 for clinically diagnosed COVID-19, or deaths with U071 or U072 as an underlying cause of death were identified.

Records with U071 or U072 as a secondary diagnosis were counted as cases if the main diagnosis was clinically related to COVID-19 (full list in online supplemental table S1). This resulted in a reduction of total cases from 2957 to 2635.

All cases were considered as severe COVID-19 since they required hospital care. Furthermore, patients were categorised into three groups with increasing severity:

- ▶ Hospitalisation due to COVID-19: n=2594.
- ▶ Intensive care due to COVID-19: n=559.
- ▶ Deaths due to COVID-19: n=184.

Register data based on Swedish hospital records have high validity.²¹ As the healthcare system was stressed during 2020 but never overloaded, it is unlikely that a significant number of potential hospital patients with COVID-19 were not admitted or included in the records.

STATISTICAL ANALYSIS

The main independent variable, CRF, was analysed in terms of three categories describing low, medium and high values of exposure. Logistic regression was used to calculate the odds for COVID-19 outcome by this exposure category, adjusted for regional examination centre, age at and year of conscription, as well as linear, quadratic and cubic terms of BMI, height, parental education, chronic disease and blood pressure at examination. Due to the rarity of events, in particular deaths from COVID-19, we used a penalised likelihood estimation (Firth method) to reduce possible small sample bias in maximum likelihood estimation.²²

Effect modification was investigated by including product terms between the main exposure (CRF categories) and other categorical predictors into the model,

overweight status (BMI ≥ 25 kg/m²), parental education, blood pressure category and current age dichotomised at median value (52 years). For each interaction analysis, the p value from an F-test comparing a model with and without interactions was assessed. Results were given in terms of ORs and 95% CIs. Muscle strength in units of newton (N) was standardised to zero mean and unit SD, and results were given in terms of OR per SD. To explore the non-linear association between muscle strength and COVID-19, we used logistic regression based on restricted cubic splines.²³ Considering the rarity of events, the ORs may be interpreted as relative risk, according to the rare disease assumption.

To investigate whether the associations with CRF or strength were explained by differences in CVD) during midlife, we conducted a mediation analysis including confounding of exposure–mediator–outcome associations with Delta method for CIs.²⁴ A supplementary analysis about the highest levels of fitness was also conducted.

Statistical analyses were performed with SAS V.9.4 (SAS Institute). Statistical significance was set at 0.05 (two-sided tests).

Ethics approval statement

The study conforms to the principles outlined in the Declaration of Helsinki. The data were anonymised before being accessed by the study authors.

RESULTS

Basic characteristics of the 1 559 187 men included in the study are shown by conscription decade in [table 1](#). The majority of deaths and hospitalisations due to COVID-19 were observed among those who underwent conscription between 1968 and 1985.

CRF in early adulthood and severity of COVID-19

Our analysis shows a protective association between higher CRF at conscription and hospitalisation, intensive care or death due to COVID-19 later in life, adjusted for examination year and place and conscript's age at examination. ORs for high versus low fitness were 0.76 (95% CI 0.67 to 0.85), 0.61 (95% CI 0.48 to 0.78) and 0.56 (95% CI 0.37 to 0.85) respectively.

This association remains statistically significant with only minor attenuation after also controlling for BMI ([figure 2](#), left panel), as well as blood pressure, height, baseline morbidity and parental education level (full results in online supplemental table S2).

In order to examine whether there was any additional or decreased protection at the highest level of CRF, categories 8 and 9 were disaggregated, but no statistically significant difference could be seen between the two subgroups. Adjusting for muscle strength did not change the results for CRF.

Interaction analyses indicated that the associations between CRF and all three COVID-19 endpoints were not modified by weight status, blood pressure, baseline

morbidity or education. For example, comparing high versus low fitness in overweight (BMI group 3–5) and non-overweight conscripts (BMI group 1–2), the association with intensive care requiring COVID-19 tended to be stronger in the overweight group, OR=0.49 (0.25–0.96), than in the non-overweight group, OR=0.65 (0.50–0.85), but not significantly so (p value for interaction=0.5). Similarly, the other three tested interactions were not statistically significant (p>0.1).

Muscle strength and severity of COVID-19

Muscular strength, a continuous variable measured in units of newton and recorded until 1997, showed associations with COVID-19 that were similar in direction to the more robust protective association with CRF. As shown in [figure 2](#), right panel, when analysed in intervals of 313 N (=1 SD), the association was statistically significant for all three outcomes, but after controlling for cardiovascular fitness, the results were attenuated (full results in online supplemental table S3). Further analysis using cubic spline analyses confirmed the association between muscle strength and COVID-19. Specifically, these analyses, illustrated in supplementary materials (online supplemental figure S1), confirmed that lower muscle strength in young adulthood has a significant linear association for all three endpoints and no significant non-linear component.

Supplementary mediation analysis

A supplementary analysis of CVD as a mediating factor in the association between CRF or muscle strength and severe COVID-19 was conducted. This analysis indicated that the direct effects of CRF and muscular strength (not mediated by CVD) are only slightly weaker than the overall total effect. Full results of this analysis are available in online supplemental table S4.

DISCUSSION

The main finding of this study is that lower CRF and lower muscle strength in young adulthood are predictive of severe illness and death due to COVID-19 15–52 years later. The protective association between cardiovascular fitness and COVID-19 severe enough to warrant hospitalisation, intensive care and/or death was not attenuated by controlling for BMI at the time of conscription examinations. It was also independent of parental education, chronic morbidity and blood pressure at conscription.

As of this date, no other comparable study has been published on the link between fitness in young adulthood and later COVID-19. Using data from the Swedish military conscript registry provided us with objective measures of fitness in a uniquely large sample. This together with the prospective design, the long and near-complete follow-up from well-validated hospital records and the death registry provides a strong ground for drawing conclusions.

Other studies point in the same direction, though with a shorter time span between exposure and outcome. Measured hand grip strength has been shown to correlate

Table 1 Description of analytical sample by conscription decade

Conscription year	1968–1975	1976–1985	1986–1995	1996–2005	All decades
N	283 082	441 414	496 484	338 207	1 559 187
Age at conscription, mean (SD)	18.5 (0.65)	18.3 (0.82)	18.3 (0.76)	18.2 (0.58)	18.3 (0.73)
Age in 2020, mean (SD)	65.9 (1.94)	57.5 (2.98)	48.0 (3.00)	38.3 (2.91)	51.8 (9.89)
CRF					
n with CRF score	248 424	380 791	358 731	155 720	1 143 670
CRF low, n (%)	89 953 (36.2)	135 569 (35.6)	85 203 (23.8)	25 546 (16.4)	336 271 (29.4)
CRF medium, n (%)	84 189 (33.9)	127 965 (33.6)	189 000 (52.7)	87 343 (56.1)	488 497 (42.7)
CRF high, n (%)	74 286 (29.9)	117 257 (30.8)	84 528 (23.6)	42 831 (27.5)	318 902 (27.9)
Muscle strength					
n with strength score in newton	277 150	435 323	449 188	n/a	1 161 661
Mean score (SD)	2 033 (299)	2 092 (315)	2 140 (315)	n/a	2 096 (313)
BMI					
Mean (SD)	21.2 (2.6)	21.7 (2.8)	22.0 (3.0)	22.7 (3.6)	21.9 (3.0)
Overweight and obese, n (%)	20 481 (1.5)	44 265 (2.2)	65 268 (3.2)	64 943 (5.6)	194 957 (3.6)
Height					
Mean cm (SD)	178.7 (6.4)	179.1 (6.5)	179.3 (6.6)	179.8 (6.7)	179.2 (6.7)
Blood pressure					
Grade 1 hypertension, n (%)	51 242 (18.1)	81 990 (18.6)	89 752 (18.5)	57 582 (21.4)	280 566 (19)
Grade 2 hypertension, n (%)	2 167 (0.8)	3 145 (0.7)	2 685 (0.6)	1 614 (0.6)	9 611 (0.7)
Baseline morbidity					
n with CVD (%)	8 025 (2.8)	10 836 (2.5)	13 836 (2.8)	4 732 (1.4)	37 429 (2.4)
n with diabetes (%)	102 (0.04)	255 (0.06)	321 (0.06)	273 (0.08)	951 (0.06)
n with kidney disease (%)	285 (0.1)	399 (0.09)	508 (0.1)	316 (0.09)	1 508 (0.1)
n with malignant cancer (%)	71 (0.03)	134 (0.03)	371 (0.07)	231 (0.07)	807 (0.05)
n with respiratory disease (%)	22 760 (8.04)	49 808 (11.3)	88 904 (17.9)	66 155 (19.6)	227 627 (14.6)
Parental education					
Low, n (%)	101 918 (52.9)	141 205 (34.8)	85 586 (17.5)	28 186 (8.4)	356 895 (25)
Medium, n (%)	77 020 (40)	213 050 (52.5)	309 246 (63.1)	222 095 (65.8)	821 411 (57.6)
High, n (%)	13 787 (7.2)	51 660 (12.7)	95 589 (19.5)	87 069 (25.8)	248 105 (17.4)
COVID-19 diagnoses					
n in hospital due to COVID-19 (%)	728 (0.26)	944 (0.21)	674 (0.14)	248 (0.07)	2 594 (0.17)
n in intensive care due to COVID-19 (%)	195 (0.07)	209 (0.05)	130 (0.03)	25 (0.01)	559 (0.04)
n dead due to COVID-19 (%)	117 (0.04)	43 (0.01)	24 (0.00)	0	184 (0.01)

CRF, cardiorespiratory fitness.

with COVID-19 hospitalisations among adults 50 years and older.¹⁰ Also being consistently inactive is strongly associated with increased risk of severe COVID-19.⁷

A possible mechanism for associations between CRF and severe COVID-19 could be higher cardiovascular morbidity (diagnosed or not) in the years after conscription examination among those with low CRF. Earlier studies using the same data have shown an association between low fitness and higher risk of heart failure, diabetes type 2, stroke, hypertension, ischaemic heart disease and psychiatric disorders.^{15 25–28} In this regard, CRF could also be a marker for a better long-term lifestyle (less smoking, better diet).

Alternatively, there could be a direct anti-inflammatory effect or a modulation of the immune system, either acquired during adolescence and early adulthood and retained during the years leading up to 2020 or continually reinforced by physical activity established earlier in life.²⁹ In a subsample of 139 609 individuals, we had access to more recent CRF data collected from a health profiling registry,³⁰ showing a clear correlation between these estimates of VO_{2max} and the fitness categories from conscription, indicating that physical fitness tracks over time.

However, only a minor part of the protective association between CRF and COVID-19 outcome was mediated by




Figure 2 Association between categories of cardiorespiratory fitness (CRF) and severe COVID-19 (left panel, n=1 143 670, hospitalisations=2006, intensive care=445 and deaths=149) and between muscle strength as a continuous predictor and severe COVID-19 (right panel, or per SD of muscle strength, n=1 161 827, hospitalisations=2252, intensive care=514 and deaths=180). All regression models were made with Firth's bias correction and adjusted for year and place, as well as age and BMI at conscription examination. BMI, body mass index.

protection from CVD in midlife. This points to a genuine effect of physical fitness acting via a modulation of the immune system. Of course, we cannot rule out a genetic predisposition to lower inflammatory levels, in conjunction with genetically higher CRF.³¹

Despite a number of strengths, the study is not without limitations, including the fact that a large number of Swedish citizens have immigrated as adults and never went through the conscription examination. Immigrants have been over-represented in the Swedish COVID-19 mortality and morbidity statistics with explanations ranging from crowded living conditions, lack of official information and occupations where working from home is not an option. This group has a lower socioeconomic status and higher degree of lifestyle-related risk factors (more obesity, more smoking and more diabetes). Importantly, this group has also lower CRF, potentially contributing to explaining their higher risk.³²

Moreover, the results from this cohort can only be generalised to men, since the conscription process was not mandatory for women. Other limitations include some protocol changes in strength and CRF measurements over the years. For instance, a rather large number of conscripts are lacking CRF scores, predominantly during the last decade where the lowest scores were not recorded. This makes it difficult to draw conclusions regarding very low CRF levels and could underestimate the adverse association. In contrast to the CRF scores, the muscular strength measurements among the conscripts are normally distributed, and the analysis shows a considerable increase in the odds of developing severe COVID-19 among those at the lowest end of the distribution. When CRF was included in the model, the effect of mus, indicating that CRF is the more important factor.

Height turned out to be independently predictive of hospitalisation with OR 1.014 (1.006–1.021) per cm (95% CI, basic model adjusted for BMI), with a possible explanation being the association between height and venous thromboembolism.³³

A related concern is the current age range of the cohort (32–70 years in 2020). All reported findings were adjusted for exact age at conscription and year of conscription examination and thus indirectly for age in 2020. To further illustrate this point, we repeated the analysis after stratification at median current age (52 years) and found only minor differences between the age groups. This underscores that protective associations of higher fitness are observed independent of cohort differences in current age or length of follow-up.

The present study has clear clinical implications. Physical fitness is a (largely) modifiable exposure and is therefore an important preventive measure both affecting later CVD and, as this study shows, severe COVID-19. The obvious public health implications ought to be to further strengthen efforts concerning fitness in the young and presumably through the life course. Anecdotal stories about severely sick athletes during 2020 (the first patient in intensive care in Italy was a marathon runner) raised the question if extremely physically fit individuals had a higher risk of severe COVID-19 but no statistically significant difference could be seen between those of very high CRF (score 9) and those with moderately-high (score 8), although the power of this analysis was low.

Looking towards the future, there are also the indirect effects of the pandemic and subsequent lockdowns on physical activity behaviours. In Sweden, a country with relatively few restrictions during the pandemic, only small changes in lifestyle habits have been shown so far,³⁴

even though the amount of time sitting down is similar to other study settings.^{35 36}

Considering the fact that women seem to run lower risk for severe COVID-19,³⁷ it is unfortunate not to be able to compare the protective effects of CRF in women and men. Using other sources of data such as the Swedish medical birth register, this might be done in a future study. Finally, although we found the results for CRF were generally robust and unaffected by adjusting for covariates such as BMI, we cannot exclude the possibility of residual confounding by unmeasured factors. This includes later exposures and comorbidities occurring between conscription and 2020.

CONCLUSION

There is growing evidence of the importance of CRF on morbidity later in life,³⁰ whereas this study shows fitness at a young age may influence the severity level of COVID-19 many years later. In the near future, updates of the COVID-19 data will make it possible to understand these effects in more detail but are unlikely to reverse the results.

The findings of the present study reinforce the need to promote regular physical activity early in life to increase the general CRF and muscular strength of the population to decrease the risk of future cardiovascular events and other conditions and to offer protection against potential consequences of future viral pandemics.

Author affiliations

¹School of Public Health and Community Medicine, Institute of Medicine, University of Gothenburg Sahlgrenska Academy, Goteborg, Sweden

²Department of Molecular and Clinical Medicine, University of Gothenburg Sahlgrenska Academy, Goteborg, Sweden

³Center for Health and Performance, University of Gothenburg, Goteborg, Västra Götaland, Sweden

⁴Department of Infectious Diseases, Institute of Biomedicine, University of Gothenburg Sahlgrenska Academy, Goteborg, Sweden

⁵Section for Clinical Neuroscience, Institute of Neuroscience and Physiology, University of Gothenburg Sahlgrenska Academy, Goteborg, Sweden

⁶Sahlgrenska University Hospital, Goteborg, Sweden

⁷Regionhälsan, Region Västra Götaland, Göteborg, Sweden

Twitter Agnes af Geijerstam @agnesafg

Contributors LL and MÅ initiated the project. AaG and KM performed all statistical analyses. AaG had main responsibility for writing the article. MB, JR, JN, MA and AR all made substantial contributions to the interpretation of the analyses, the structure and content of the manuscript and have read and approved of the final draft. All authors have agreed to be accountable for all aspects of the work.

Funding This work was supported by the EpiLife-Teens Research Program (FORMAS2012-00038), the Swedish ALF-agreement (ALFGBG-720201) and the Swedish Research Council (02508, VRREG 2019-00193, 2020-05792).

Disclaimer The funding sources had no role in study design, collection, analysis and interpretation of data, the writing of the report, or the decision to submit the article for publication.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval The Ethics Committee of the University of Gothenburg and Confidentiality Clearance at Statistics Sweden approved the study (EPN Reference numbers EPN 462–14 and 567–15; T174-15, T653-17, T196-17, T 2020–01325,

T 2020–02420). The requirement for informed consent was waived by the Ethics Committee of the University of Gothenburg for secondary analysis of existing data.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data may be obtained from a third party and are not publicly available. The data used in this study is available on request from the Swedish National Board of Health and Welfare, the Swedish intensive care registry and Statistics Sweden.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Agnes af Geijerstam <http://orcid.org/0000-0002-0897-6548>

Jenny Nyberg <http://orcid.org/0000-0002-4336-3886>

REFERENCES

- Clark A, Jit M, Warren-Gash C, *et al*. Global, regional, and national estimates of the population at increased risk of severe COVID-19 due to underlying health conditions in 2020: a modelling study. *Lancet Glob Health* 2020;8:e1003–17.
- Hamer M, Gale CR, Kivimäki M, *et al*. Overweight, obesity, and risk of hospitalization for COVID-19: a community-based cohort study of adults in the United Kingdom. *Proc Natl Acad Sci U S A* 2020;117:21011–3.
- O'Driscoll M, Ribeiro Dos Santos G, Wang L, *et al*. Age-Specific mortality and immunity patterns of SARS-CoV-2. *Nature* 2021;590:140–5.
- Ross R, Blair S, Arena R. American heart association physical activity Committee of the Council on lifestyle and cardiometabolic health; Council on clinical cardiology; Council on epidemiology and prevention; Council on cardiovascular and stroke nursing; Council on functional genomics and translational biology; stroke Council. Importance of assessing cardiorespiratory fitness in clinical practice: a case for fitness as a clinical vital sign: a scientific statement from the American heart association. *Circulation* 2016;134:e653–99.
- Zbinden-Foncea H, Francaux M, Deldicque L, *et al*. Does high cardiorespiratory fitness confer some protection against proinflammatory responses after infection by SARS-CoV-2? *Obesity* 2020;28:1378–81.
- Yates T, Razieh C, Zaccardi F, *et al*. Obesity, walking PACE and risk of severe COVID-19 and mortality: analysis of UK Biobank. *Int J Obes* 2021;45:1155–9.
- Sallis R, Young DR, Tartof SY, *et al*. Physical inactivity is associated with a higher risk for severe COVID-19 outcomes: a study in 48 440 adult patients. *Br J Sports Med* 2021. doi:10.1136/bjsports-2021-104080. [Epub ahead of print: 13 Apr 2021].
- Salgado-Aranda R, Pérez-Castellano N, Núñez-Gil I, *et al*. Influence of baseline physical activity as a modifying factor on COVID-19 mortality: a single-center, retrospective study. *Infect Dis Ther* 2021;10:801–14.
- In, Brawner CA, Ehrman JK, Bole S, *et al*. Inverse relationship of maximal exercise capacity to hospitalization secondary to coronavirus disease 2019. *Mayo Clin Proc* 2021;96:32–9.
- Cheval B, Sieber S, Maltagliati S. Muscle strength is associated with COVID-19 hospitalization in adults 50 years of age and older. *MedRxiv* 2021.
- Aberg MAI, Pedersen NL, Torén K, *et al*. Cardiovascular fitness is associated with cognition in young adulthood. *Proc Natl Acad Sci U S A* 2009;106:20906–11.
- Gao M, Piernas C, Astbury NM. Associations between body-mass index and COVID-19 severity in 6·9 million people in England: a



- prospective, community-based, cohort study. *Lancet Diabetes Endocrinol* 2021.
- 13 Prince SA, Adamo KB, Hamel ME, *et al*. A comparison of direct versus self-report measures for assessing physical activity in adults: a systematic review. *Int J Behav Nutr Phys Act* 2008;5:56–24.
 - 14 Berg C, Rosengren A, Aires N, *et al*. Trends in overweight and obesity from 1985 to 2002 in Göteborg, West Sweden. *Int J Obes* 2005;29:916–24.
 - 15 Åberg ND, Kuhn HG, Nyberg J, *et al*. Influence of cardiovascular fitness and muscle strength in early adulthood on long-term risk of stroke in Swedish men. *Stroke* 2015;46:1769–76.
 - 16 Nordesjö L, Schéle R. Validity of an ergometer cycle test and measures of isometric muscle strength when prediction some aspects of military performance. *Swed J Def Med* 1974;10:11–23.
 - 17 Lissner L, Mehlig K, Rosengren A, *et al*. A growing social divide in body mass index, strength, and fitness of Swedish male conscripts. *J Adolesc Health* 2019;65:232–8.
 - 18 World Health Organization. The Asia-Pacific perspective: redefining obesity and its treatment 2000.
 - 19 Cole TJ, Lobstein T. Extended International (IOTF) body mass index cut-offs for thinness, overweight and obesity. *Pediatr Obes* 2012;7:284–94.
 - 20 Williams B, Mancia G, Spiering W. ESC/ESH guidelines for the management of arterial hypertension: the task force for the management of arterial hypertension of the European Society of cardiology (ESC) and the European Society of hypertension (ESH). *Eur Heart J* 2018;39:3021–104.
 - 21 Ludvigsson JF, Andersson E, Ekblom A, *et al*. External review and validation of the Swedish national inpatient register. *BMC Public Health* 2011;11:1–16.
 - 22 King G, Zeng L. Logistic regression in rare events data. *Polit. anal.* 2001;9:137–63.
 - 23 Li R, Hertzmark E, Louie M. The SAS lgtphcurv9 macro. *Boston Channing Lab* 2011.
 - 24 Valeri L, Vanderweele TJ. Mediation analysis allowing for exposure-mediator interactions and causal interpretation: theoretical assumptions and implementation with SAS and SPSS macros. *Psychol Methods* 2013;18:137–50.
 - 25 Lindgren M, Åberg M, Schaufelberger M, *et al*. Cardiorespiratory fitness and muscle strength in late adolescence and long-term risk of early heart failure in Swedish men. *Eur J Prev Cardiol* 2017;24:876–84.
 - 26 Crump C, Sundquist J, Winkleby MA, *et al*. Interactive effects of obesity and physical fitness on risk of ischemic heart disease. *Int J Obes* 2017;41:255–61.
 - 27 Crump C, Sundquist J, Winkleby MA, *et al*. Interactive effects of physical fitness and body mass index on the risk of hypertension. *JAMA Intern Med* 2016;176:210–6.
 - 28 Crump C, Sundquist J, Winkleby MA, *et al*. Physical fitness among Swedish military conscripts and long-term risk for type 2 diabetes mellitus: a cohort study. *Ann Intern Med* 2016;164:577–84.
 - 29 Gleeson M, Bishop NC, Stensel DJ, *et al*. The anti-inflammatory effects of exercise: mechanisms and implications for the prevention and treatment of disease. *Nat Rev Immunol* 2011;11:607–15.
 - 30 Ekblom-Bak E, Ekblom B, Söderling J, *et al*. Sex- and age-specific associations between cardiorespiratory fitness, CVD morbidity and all-cause mortality in 266,109 adults. *Prev Med* 2019;127:105799.
 - 31 Burtscher J, Millet GP, Burtscher M. Low cardiorespiratory and mitochondrial fitness as risk factors in viral infections: implications for COVID-19 2020.
 - 32 Lindgren M, Börjesson M, Ekblom Örgan, *et al*. Physical activity pattern, cardiorespiratory fitness, and socioeconomic status in the SCAPIS pilot trial - A cross-sectional study. *Prev Med Rep* 2016;4:44–9.
 - 33 Braekkan SK, Borch KH, Mathiesen EB, *et al*. Body height and risk of venous thromboembolism: the Tromsø study. *Am J Epidemiol* 2010;171:1109–15.
 - 34 Blom V, Lönn A, Ekblom B, *et al*. Lifestyle habits and mental health in light of the two COVID-19 pandemic waves in Sweden, 2020. *Int J Environ Res Public Health* 2021;18:3313.
 - 35 Cheval B, Sivaramakrishnan H, Maltagliati S, *et al*. Relationships between changes in self-reported physical activity, sedentary behaviour and health during the coronavirus (COVID-19) pandemic in France and Switzerland. *J Sports Sci* 2021;39:699–704.
 - 36 Flanagan EW, Beyl RA, Fearnbach SN, *et al*. The impact of COVID-19 Stay-At-Home orders on health behaviors in adults. *Obesity* 2021;29:438–45.
 - 37 Peckham H, de Grujter NM, Raine C, *et al*. Male sex identified by global COVID-19 meta-analysis as a risk factor for death and ICU admission. *Nat Commun* 2020;11:1–10.

Supplementary material

Table S1. Included main diagnoses divided by categories when COVID-19 is secondary diagnosis.

Included categories	ICD-codes
COVID-related symptoms	
Cough	R05
Abnormalities of breathing	R06
Pain in throat and chest	R07
Other symptoms and signs involving the circulatory and respiratory system	R09
Dizziness and giddiness	R42
Fever of other or unknown origin	R50
Headache	R51
Malaise and fatigue	R53
Syncope and collapse	R55
Upper and lower respiratory tract infections	
Acute nasopharyngitis	J00
Acute tonsillitis	J03
Acute upper respiratory infections of multiple and unspecified sites	J06
Influenza due to other identified influenza virus	J10
Other viral pneumonia	J128
Viral pneumonia, unspecified	J129
Pneumonia due to <i>Streptococcus pneumoniae</i>	J13
Bacterial pneumonia, not elsewhere classified	J15
Pneumonia due to other specified infectious organisms	J168
Pneumonia in diseases classified elsewhere	J17
Pneumonia, unspecified organism	J18
Unspecified acute lower respiratory infection	J22
Coronavirus infection, unspecified	B342
Other viral infections of unspecified site	B348
Viral infection, unspecified	B349
Coronavirus as the cause of diseases classified elsewhere	B972
Other and unspecified infectious diseases	B99
Respiratory disorders	
Pulmonary embolism	I26
Acute respiratory distress syndrome	J80
Pulmonary edema	J81
Pleural effusion not elsewhere classified	J90
Respiratory failure, not elsewhere classified	J96
Respiratory disorders in diseases classified elsewhere	J99
Obstructive Airway Diseases	
Acute bronchitis due to other specified organisms	J208
Acute bronchitis, unspecified	J209
Acute bronchiolitis due to other specified organisms	J218
Acute bronchiolitis, unspecified	J219
Other chronic obstructive pulmonary disease	J44
Asthma	J45
Cardiac diseases	
Viral carditis	B332
Chronic ischemic heart disease	I25
Acute pericarditis	I30
Pericarditis in diseases classified elsewhere	I32
Acute myocarditis, unspecified	I40
Myocarditis in diseases classified elsewhere	I41
Atrial fibrillation and flutter	I48

Heart failure	I50
Abnormalities of heart beat	R00
Kidney disorders	
Acute kidney failure	N17
Chronic kidney disease	N18
Unspecified kidney failure	N19
Electrolyte disorders	
Other disorders of fluid, electrolyte and acid-base balance	E87

Table S2: Results of main analysis. Association between conscript's fitness at time of examination and hospitalization, intensive care and death due to COVID-19 (OR, 95% CI, Firth's bias correction).

Fitness level	Basic model ^a	Basic model controlled for BMI ^b	Basic model controlled for blood pressure n=1 140 514	Basic model controlled for baseline morbidity ^c n=1 143 670	Basic model controlled for BMI ² and height N=1 143 670	Basic model controlled for parental education n=1 031 246	Basic model controlled for BMI, height, blood pressure, morbidity and parental education n=1 028 122
	n=1 143 670	n=1 143 670					
All hospitalization due to COVID-19.							
	cases=2 006	cases=2 006	cases=2 003	cases=2 006	cases=2 006	cases=1 710	cases=1 707
Low (1-5)	1	1	1	1	1	1	1
Medium (6-7)	0.91 (0.82-1.01)	0.92 (0.83-1.01)	0.91 (0.82-1.01)	0.91 (0.82-1.01)	0.91 (0.82-1.01)	0.91 (0.82-1.02)	0.91 (0.81-1.02)
High (8-9)	0.76 (0.67-0.85)	0.73 (0.65-0.83)	0.76 (0.68-0.85)	0.76 (0.67-0.85)	0.71 (0.63-0.81)	0.74 (0.65-0.85)	0.73 (0.64-0.83)
AUROC	0.66	0.66	0.66	0.66	0.66	0.67	0.67
Intensive care due to COVID-19							
	cases=445	cases=445	cases=445	cases=445	cases=445	cases=363	cases=363
Low (1-5)	1	1	1	1	1	1	1
Medium (6-7)	0.76 (0.62-0.95)	0.77 (0.62-0.95)	0.76 (0.62-0.95)	0.77 (0.62-0.95)	0.76 (0.61-0.94)	0.76 (0.60-0.96)	0.76 (0.60-0.96)
High (8-9)	0.61 (0.48-0.78)	0.58 (0.45-0.75)	0.61 (0.48-0.78)	0.62 (0.48-0.79)	0.57 (0.44-0.74)	0.60 (0.45-0.80)	0.60 (0.46-0.80)
AUROC	0.69	0.71	0.70	0.69	0.71	0.71	0.71
Mortality due to COVID-19							
	cases=149	cases=149	cases=149	cases=149	cases=149	cases= 111	cases=111
Low (1-5)	1	1	1	1	1	1	1
Medium (6-7)	0.67 (0.46-0.96)	0.64 (0.44-0.94)	0.67 (0.47-0.97)	0.66 (0.46-0.96)	0.63 (0.43-0.92)	0.53 (0.34-0.83)	0.53 (0.34-0.81)
High (8-9)	0.56 (0.37-0.85)	0.52 (0.34-0.80)	0.57 (0.38-0.87)	0.56 (0.37-0.85)	0.50 (0.32-0.77)	0.52 (0.32-0.83)	0.50 (0.31-0.81)
AUROC	0.81	0.82	0.81	0.81	0.82	0.82	0.82

^a adjusted for year and place of examination, and age at examination.

^b BMI, BMI², BMI³

^c including CVD, diabetes, kidney disease, respiratory disease and cancer.

Table S3: Association between muscle strength at conscription and severity of COVID disease (OR for disease by 1 SD higher muscle strength (SD = 313 N), 95% CI, Firth's bias correction)

Outcome	Total n	Cases	OR (95% CI)	AUROC
<i>Hospital^a</i>	1 161 914	2 252	0.89 (0.84-0.93)	0.64
<i>ICU^a</i>		514	0.88 (0.79-0.97)	0.68
<i>Mortality^a</i>		180	0.83 (0.70-0.99)	0.80
<i>Hospital^b</i>	952 225	1 850	0.90 (0.86-0.95)	0.64
<i>ICU^b</i>		421	0.90 (0.80-1.01)	0.68
<i>Mortality^b</i>		146	0.85 (0.70-1.03)	0.79

^a basic model adjusted for BMI, BMI², BMI³ and height.

^b above model further adjusted for fitness categories.

Figure S1: Results of restricted cubic spline analyses of continuous muscle strength and severity of COVID-19 adjusted for age, year, and place of conscript examination, BMI, BMI², BMI³, and height (total number of observations = 1 161 914).




Table S4. Results of mediation analysis between cardiovascular fitness, muscle strength, cardiovascular disease and severe COVID-19 (OR with 95%CI).

Outcome	Direct	Indirect	Total	Proportion-mediated ^c
Muscle strength^a				
Hospital	0.893 (0.850-0.937)	0.995 (0.994-0.996)	0.888 (0.846-0.933)	4.0 %
ICU	0.880 (0.796-0.973)	0.997 (0.995-0.998)	0.877 (0.793-0.970)	2.2 %
Mortality	0.855 (0.720-1.015)	0.992 (0.989-0.995)	0.848 (0.714-1.007)	4.5 %
CRF^{a,b}				
Hospital ^a	0.860 (0.810-0.913)	0.987 (0.985-0.989)	0.849 (0.800-0.901)	7.4 %
ICU ^a	0.772 (0.679-0.877)	0.991 (0.987-0.995)	0.765 (0.673-0.869)	3.0 %
Mortality ^a	0.725 (0.578-0.909)	0.978 (0.969-0.986)	0.709 (0.566-0.889)	5.5 %

^a adjusted for age, year, and place of conscript examination, BMI, BMI², BMI³ and height.

^b treating the 3-level variable for CRF as ordinal score.

^c calculated from direct and indirect effect odds ratios as $OR^{direct} (OR^{indirect} - 1) / (OR^{direct} OR^{indirect} - 1)$.

VanderWeele T, Mediation Analysis: A Practioner's Guide, Annu. Rev. Public Health 2016.37:17-32